
MODERN INTRODUCTORY ANALYSIS

DOLCIANI

BECKENBACH: JURGENSEN

ONNELLY WOOTON



ABOUT THE AUTHORS

Mary P. Dolciani, Associate Dean for Academic Services and Professor, Department of Mathematics, Hunter College of the City University of New York. Dr. Dolciani has been a director and teacher in numerous National Science Foundation and New York State Education Department institutes for mathematics teachers, and Visiting Secondary School Lecturer for the Mathematical Association of America.

Edwin F. Beckenbach, Professor of Mathematics, University of California, Los Angeles. Dr. Beckenbach has been a team member of National Council of Teachers of Mathematics summer writing projects. He is Chairman of the Committee on Publications of the Mathematical Association of America.

Alfred J. Donnelly, Master Instructor and holder of the William Pitt Oakes Chair of Mathematics, Culver Military Academy and Culver Academy for Girls. Mr. Donnelly brings a rich background of both mathematics study and teaching to his authorship.

Ray C. Jurgensen, Chairman of Mathematics Department and holder of the Eppley Chair of Mathematics, Culver Military Academy and Culver Academy for Girls, Culver, Indiana. Mr. Jurgensen is a regular reader for the Advanced Placement Examinations of the College Entrance Examination Board.

William Wooton, Mathematics Consultant, Vista, California, Unified School District. Mr. Wooton was formerly Professor of Mathematics at Los Angeles Pierce College and has been a teacher at the junior and senior high school levels.

EDITORIAL ADVISER

Albert E. Meder, Jr., Professor of Mathematics and Dean of the University, Emeritus, Rutgers University, the State University of New Jersey.

COPYRIGHT @ 1974, 1970, 1967, 1964 BY HOUGHTON MIFFLIN COMPANY

All rights reserved. No part of this work may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying and recording, or by any information storage or retrieval system, without permission in writing from the publisher. Printed in the U.S.A.

Student Edition ISBN: 0-395-18420-7 Teacher's Edition ISBN: 0-395-18418-5

Statements and Sets

in Mathematics

One way to view mathematics is to think of it as a study of patterns. In this book, you will learn to work with a wide variety of patterns, some familiar to you, such as $x^2 - y^2 = (x + y)(x - y)$, and others quite new. In order to learn the mathematics you will encounter, you will need to understand the logical machinery used to discuss mathematical statements; this is what you will study in the present chapter.

SIMPLE STATEMENTS AND SETS

1-1 Logical Statements; Sets

In studying algebra and geometry you have discovered many logical connections between mathematical statements. But do you know what constitutes a statement in logic?

A statement or proposition is a set of symbols (and words are symbols, of course) which forms a meaningful assertion that is either true or false, but not both true and false.

The following assertions are examples of statements:

True Statements

False Statements

(1) 2 + 3 = 5.

(1) 0 > 1.

- (2) Shakespeare wrote Hamlet.
- (2) Plato was a Chinese philosopher.
- Ottawa is not a province of Canada.
- (3) $\sqrt[3]{125} \neq 5$.

On the other hand, none of the following sentences is a statement because none can be said to be definitely true or definitely false.

- (1) Give me liberty or give me death.
- (2) $x^2 + 1 = 5$.
- (3) This sentence is false.

2 CHAPTER ONE

Logical statements in mathematics frequently involve the use of symbols of equality and inequality. You should recall that the symbol "=" represents the word "equals" or the phrase "is equal to," and is used between expressions to show that they represent the same object. Similarly, the inequality symbols ">," "<," "≥," and "≤" represent "is greater than," "is less than," "is greater than or equal to," and "is less than or equal to," respectively. These symbols are used in sentences and statements comparing numbers. A bar, / or |, is used in conjunction with certain symbols to denote negation. For example, the symbol "≠" placed between two expressions asserts that the expressions do not represent the same object, and the symbol < stands for "is not less than."

Some mathematical statements concern membership in a set. You should recall from your earlier study of mathematics that the objects in a set are called members or elements of the set, and are said to belong to or to be contained in the set. The symbol \in is used to denote "is an element of," while \notin means "is not an element of." For example, if J is the set of integers, then $3 \in J$ is a true statement, whereas $4 \notin J$ is a false statement.

A statement that two sets are equal is an assertion that they contain the same elements. Thus, the statement A = B means that A and B name the same set. This definition of the equality of sets implies that to specify a set, you must be able to identify its elements. You can sometimes specify a set by listing the names of its elements within braces, $\{ \}$. For example, $\{-1,0,1\}$, read "the set whose members are -1,0, and 1," is a roster (list) of the numbers -1,0, and 1. Quite frequently, however, a set has so many elements that it is inconvenient or impossible to list them all. Thus, we may write $\{1,2,3,\ldots,50\}$ and expect the reader to understand that the set specified contains the integers from 1 to 50, inclusive. Similarly, $\{1,2,3,\ldots\}$ specifies the set of all positive integers.

The members of a set can, themselves, be sets. $\{\{1,3\}, \{5,7\}, \{9,11\}\}$, for example, is the set whose members are $\{1,3\}, \{5,7\}$, and $\{9,11\}$. Observe that the set $\{\{a,b\}\}$ is not the same as the set $\{a,b\}$. The first contains just one element, $\{a,b\}$, while the second contains two elements, a and b.

Another way to specify a set consists in giving a rule or condition that enables you to decide whether or not any given object belongs to the set.

For example, {the teachers of mathematics}, read "the set of the teachers of mathematics," specifies a set whose roster contains many thousands of names. Of course, you certainly recognize that every mathematics teacher you know belongs to




Figure 1-1

{the teachers of mathematics}. Whenever each element of a set R is also an element of a set S, we say that R is a subset of S, in symbols " $R \subset S$." Thus, {mathematics teachers you know} \subset {teachers of mathematics}. Diagrams such as the one shown are used to illustrate a set and a subset; they are called Venn diagrams.

Since every positive integer is a member of the set of positive integers, it is certainly true that the set of positive integers is a subset of itself. Each set is said to be the improper subset of itself; every other subset is called a proper subset of the set.

Can you list the members of the set of integers between $\frac{1}{2}$ and $\frac{2}{3}$? This set contains no elements at all and is therefore called the **empty set** or **null set**. We use the symbol \emptyset , written without braces, to designate the empty set. Because of our agreement on the meaning of equality of sets, there is only one empty set, \emptyset . Furthermore, \emptyset is taken to be a *proper* subset of every set except itself.

Exercises

Which of the sentences in Exercises 1–16 are statements? Of the statements, which are true? Give a reason for each answer.

A

1.
$$5 \cdot 4 = 20$$

2.
$$5-4=2$$

$$p$$
 3. 3 + 7(2) = 2 + 15

4.
$$7(4+2) = 7 \cdot 4 + 7 \cdot 2$$

$$\frac{8}{2}$$
 5. $\frac{6+2}{2}$ = 6 + 1

6. $(-2)^3 \ge (-2)^4$

7.
$$1+3 \neq 1+6$$

8.
$$2+3>0$$

9. Man the lifeboats!

In Exercises 11–16, let $J = \{\text{integers}\} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$.

11.
$$\{1, 2, 7\} \subset J$$

12. 0 ∈ J

14. ∅ is not a subset of J.

15.
$$-1 \subset J$$

Copy each sentence, making it a true statement by replacing each question mark with a numeral or with one of the symbols =, $\not=$, \in , \notin , \subset . (Note: There may be more than one correct answer.)

17.
$$-3 \times ? = -3$$

$$\sqrt{18}$$
 18. $7 \times 7 = 0$

19.
$$5(? - 3) = 20$$

20.
$$\frac{24-6}{6}$$
? $4-6$

21.
$$\frac{10 \times 4}{2}$$
 ? 5 × 2

22.
$$\frac{21+?}{3} \neq 7+1$$

23.
$$\{0\cdot 2\} \subset \{1, 3, ?\}$$

24.
$$\{1, \frac{6}{2}, -1\} = \{-2 \div 2, 3, ? \times 4\}$$

25. {0, 2, 4} ? {the even integers}.

27.
$$\frac{8 \times ?}{4} \in \{8\}$$
 29. 0 ? $\{0\}$ **31.** $\{1, \{2\}\}$? $\{\{2\}, 1\}$

32. {1, {2}} ? {{1}, 2}

Let $U = \{-4, 0, 8\}$. List all the subsets of U that

- 33. Have exactly one element. 35. Have at least two elements.
- 34. Have no elements.
- 36. Have no more than 2 elements.

Which of the following rosters or rules (Exercises 37-40) specify sets? Justify your answer by explaining whether or not it can be decided that an arbitrary object is or is not a member.

- A 37. {the authors of this textbook}
 - 38. {bell, book, candle}
- £ 39. (the digits appearing in the decimal numeral for 1/2)
 - 40. {3, 1, 4, 5, 9, 2, 6, . . . }
- Make a list of all subsets of {∅, {∅}}.
 - 42. A set contains n > 0 elements. How many subsets does it have?
 - 43. A set containing k+1 elements has 8 more subsets than a set containing k elements. Find k.
 - Argue that if A ⊂ B and B ⊂ A, then A = B.
 - Argue that if A ⊂ B and B ⊂ C, then A ⊂ C.
 - 46. Can an element of a set be a subset of the set? Justify your answer.

ANSWERS FOR ODD-NUMBERED EXERCISES

3

Section 1-1, pages 3-4 1. true statement 3. true statement 5. false statement 7. true statement 9, not a statement 11, true statement 13, false statement 15, false statement 17, 1 19, 7 21, per 4 23. 0 25. C or ∉ or ≠ 27. 4 29. € or ≠ 31. = or C 33. [-4], [0], [8] 35. [-4,0], [0,8], $\{-4, 8\}, \{-4, 0, 8\}$ 37. set 39. set 41. $\{\emptyset\}, \{\{\emptyset\}\}, \{\emptyset, \{\emptyset\}\}, \emptyset$ 43. k = 3

ANSWERS FOR ODD NUMBERED EXERCISES Section 1-1, pages 3-4

- 1. true statement
- 7. true statement
- 13. false statement
- 19. 7
- 25. ⊂ or ∉ or ≠
- $31. = \text{or} \subset$
- 37. set
- 43. k = 3

- 3. true statement
- 9. not a statement
- 15. false statement
- 21. ≠ or €
- 27. 4
- 33. {-4}, {0}, {8}
- 39. set

- 5. false statement
- 11. true statement
- 17. 1
- 23. 0
- 29. ∈ or ≠
- 35. {-4,0}, {0,8}, {-4,8}, {-4,0,8}
- 41. {ø}, {{Ø}}, {Ø, {Ø}}